

Reply to Cheng, Harko, and Shabad

Jes Madsen

Department of Physics and Astronomy, Aarhus University, DK-8000 Århus C, Denmark

(Dated: October 14, 2008)

As demonstrated in [1] there is a universal relation for the maximal electric charge of any spherical object with radius $R \gg m_e^{-1} \approx 4 \times 10^2$ fm,

$$Z_\infty(R) \approx 2m_e R / \alpha = 0.71 R_{\text{fm}}, \quad (1)$$

where Z_∞ is the maximal charge possible with infinite time available for electron-positron pair creation in a supercritical field, m_e is the electron mass, α is the fine structure constant, and R_{fm} is the radius in fm. For $R \approx m_e^{-1}$ there is a smooth transition (denoted $Z^{\text{Mads}}(R)$ in [2]) from the object-specific core charge without electrons, $Z_{\text{core}}(R)$, to the universal asymptotic relation $Z_\infty(R)$. The universal relation changes for $R \gg 25m_e^{-1} \approx 10^4$ fm into $Z(R) \approx 11.2m_e^2 R^2 = 7 \times 10^{-5} R_{\text{fm}}^2$, if only a finite time is available for pair creation [1].

The authors of [2] consider only the infinite-time case. They argue that $Z^{\text{Mads}}(R)$ is only one of two possible boundaries in the charge-radius plane that act as attractors of charged objects, the other being the threshold charge $Z_{\text{thr}}^{(1)}(R)$ where the electric charge is just sufficient for the first possible pair creation to take place. The charge-radius plane should then be divided into three regimes. (i) $Z_{\text{core}}(R) > Z^{\text{Mads}}(R)$, where pair creation saturates the Pauli principle with electrons partly shielding the core charge and reducing the net charge of the sphere to $Z^{\text{Mads}}(R)$ as found in [1]; (ii) $Z_{\text{thr}}^{(1)}(R) < Z_{\text{core}}(R) < Z^{\text{Mads}}(R)$ [3], where “the body creates pairs till the net charge diminishes down to $Z_{\text{thr}}^{(1)}(R)$ ” [2]; (iii) $Z_{\text{thr}}^{(1)}(R) > Z_{\text{core}}(R)$, where no pair creation takes place.




FIG. 1: $Z(R)$ -curves taken from [1], [2] and [4].

Ref. [2] calculates $Z_{\text{thr}}^{(n)}(R)$, where n enumerates s-wave solutions that dive into the negative energy continuum below $-m_e$, by solving the single particle Dirac equation in a large radius approximation for a spherical shell core potential and finds that $Z_{\text{thr}}^{(n)}(R) \rightarrow Z_\infty(R)$ for $R \rightarrow \infty$. This asymptotic behavior was previously demonstrated for s- and p-waves for a shell core charge in [4] and there taken to confirm the universal relation of [1].

The idea [2] that $Z_{\text{thr}}^{(1)}(R)$ should act as an attractor in the charge-radius plane is based on a misinterpretation of the solutions to the Dirac equation diving into the negative energy continuum below $-m_e$. In particular, Eqs. (1) and (2) in [2] are not applicable to the self-consistent problem considered in [1] since the screening charge from the electrons formed is not included. A detailed discussion of the underlying physics is given in [5]. When $Z_{\text{thr}}^{(2)}(R) > Z_{\text{core}}(R) > Z_{\text{thr}}^{(1)}(R)$ ((1) and (2) denote the two lowest-lying solutions found in [4], not the s-waves from [2]) only the $1s_{1/2}$ -state is diving into the negative energy continuum. This shields the core charge by two units. For $Z_{\text{thr}}^{(3)}(R) > Z_{\text{core}}(R) > Z_{\text{thr}}^{(2)}(R)$ additional screening by $p_{1/2}$ electrons becomes available, etc. As the number of electrons increases, one must self-consistently include their charge [5], [6]. For $Z_{\text{core}}(R) > Z^{\text{Mads}}(R)$ all available Fermi levels are filled and the relation of [1] is followed by any sphere with $Z_{\text{core}}(R) > Z^{\text{Mads}}(R)$. All objects with $Z^{\text{Mads}}(R) > Z_{\text{core}}(R) > Z_{\text{thr}}^{(1)}(R)$ are partly screened, and a self-consistent analysis could map out a sequence of $Z(R)$ curves populated in steady state. Contrary to the claim of [2], $Z_{\text{thr}}^{(1)}(R)$ does not act like an attractor but marks the onset of pair production.

I thank Berndt Müller, Constantin Klier, and in particular Johann Rafelski for useful comments.

- [1] J. Madsen, Phys. Rev. Lett. **100**, 151102 (2008).
- [2] K.S. Cheng, T. Harko, and A.E. Shabad, arXiv:0809.2310.
- [3] [2] states the opposite inequality; presumably a misprint.
- [4] D.A. Dicus, W.W. Repko, and V.L. Teplitz, arXiv:0808.1739. A slightly lower asymptotic charge was found for a uniform core charge because the potential was kept fixed regardless of the electrons. In [1] any core charge is screened into a shell-like charge, so the universal $Z_\infty(R)$ is reached for all initial charge distributions.
- [5] W. Greiner, B. Müller, and J. Rafelski, *Quantum Electrodynamics of Strong Fields* (Springer-Verlag, Berlin, 1985).
- [6] B. Müller and J. Rafelski, Phys. Rev. Lett. **34**, 349 (1975).

